

CORRECTION

Open Access

Correction: ingestion of a single serving of saury alters postprandial levels of plasma n-3 polyunsaturated fatty acids and long-chain monounsaturated fatty acids in healthy human adults

Zhi-Hong Yang*, Hiroko Miyahara, Jiro Takeo and Masashi Katayama

After publication of this manuscript [1], we noted that one of the monounsaturated fatty acid isomers had been mislabelled in the results, and in Figure 2A and Table 2.

In the Results and discussion, the sentence, “Concerning plasma MUFA levels, long-chain MUFA C20:1 (n-9 and n-7) and C22:1 (n-11 and n-9) peaked at 2 hr post-ingestion (Figure 2a and b)” should read, “Concerning plasma MUFA levels, long-chain MUFA C20:1 (n-11, n-9 and n-7) and C22:1 (n-11 and n-9) peaked at 2 hr post-ingestion (Figure 2a and b)”.

The figure and table have also been corrected (Figure 1, Table 1).

We apologise for this error, which was due to a misreading of the gas chromatography reading.

* Correspondence: yangzh@nissui.co.jp

Central Research Laboratory, Tokyo Innovation Center, Nippon Suisan Kaisha, Ltd., 32-3 Nanakuni 1 Chome Hachioji, Tokyo 192-0991, Japan




Figure 1 Changes in plasma levels of MUFA with aliphatic tails >18C atoms. Plasma levels (% of total plasma fatty acids) of C20:1 (n-11, n-9 and n-7) (A), C22:1 C20:1 n-9 (B), and total long-chain MUFA (C20:1 and C22:1 isomers combined) (C) are shown. The data were collected before and after the ingestion of a single saury meal. Values represent the mean \pm SE, n = 5. * P < 0.05; ** P < 0.01; *** P < 0.001 as compared to pre-ingestion values (time 0).

Table 1 Major fatty acid composition of saury used in the study, correction shown in red

FA (%)	
C14:0	6.22
C16:0	10.37
C18:0	1.80
C20:0	0.17
Total saturated FA	18.56
C16:1 n-7	1.94
C18:1 n-9	4.07
C20:1 n-11	12.55
C20:1 n-9	4.15
C22:1 n-11	20.08
C22:1 n-9	1.22
Total MUFA	44.01
C18:2n-6	1.32
C18:3n-6	0.16
C20:2n-6	0.26
C20:4n-6	0.42
Total n-6 PUFA	2.16
C18:3n-3	1.27
C20:3n-3	0.18
C20:5n-3	5.24
C22:5n-3	1.09
C22:6n-3	12.07
Total n-3 PUFA	19.85
n-3/n-6 PUFA ratio	9.19

FA: Fatty acids, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids.

Received: 1 November 2012 Accepted: 1 November 2012
 Published: 8 November 2012

Reference

1. Yang ZH, Miyahara H, Takeo J, Katayama M: Ingestion of a single serving of saury alters postprandial levels of plasma n-3 polyunsaturated fatty acids and long-chain monounsaturated fatty acids in healthy human adults. *Lipids Health Dis* 2012, **11**(1):95.

doi:10.1186/1476-511X-11-151

Cite this article as: Yang et al.: Correction: ingestion of a single serving of saury alters postprandial levels of plasma n-3 polyunsaturated fatty acids and long-chain monounsaturated fatty acids in healthy human adults. *Lipids in Health and Disease* 2012 **11**:151.